

The acceleration of charged particles in plasma at the separation of charges and generation of electrical domains

A. G. Oreshko

*Moscow Aviation Institute (National Aerospace Research University), Volokolamskoye shosse,
125871 Moscow, Russia
Oreshko_Alex@mail.ru*

The main aim of this work was to establish the mechanism of acceleration of charged particles in the plasma. In work the direct proofs of the mechanism of charged particles acceleration in the plasma at the separation of charges and generation of electrical domain were obtained and the theoretical calculations were carried. As a result of the interaction of transversal electromagnetic waves and particles, the particles gain energy. The value of acquired energy depends on the values of intensity and induction of fields and time of particles interaction with the wave.

It is known that nuclear fusion in the Sun is realized by using the accelerating mechanism [1]. The mechanism of charged particles acceleration in plasma is associated with generation in a plasma of electrical domains. The electric domains were first detected in solid-state semiconductor plasma [2]. Later electric domains were detected in the plasma glow discharge [3] and in the cathode plasma in a magnetically insulated diode of the accelerator electron [4]. Here and hereinafter under the electric domain implied the quasi-neutral in whole system which consists of a region (layer) with surplus negative charge and a region (layers) with surplus positive charge. The distance between these two regions (layer) of the domain exceeds the Debye screening length. Generation of the domain in plasma current tube is equivalent to the transformation of an element with an ohmic character of the impedance in the element with capacitive character. The charges on the plasma “capacitor” plates (layers) are defined by the next expression [4]

$$q(t) = q_0 \exp(-t/\tau) \quad (1)$$

in which τ is characteristic time of charges separation. It determined by the Maxwell relaxation time [5]

$$\tau_M = \frac{\varepsilon}{4 \pi e n \mu_e} \quad (2)$$

where ε is a dielectric permittivity and μ_e is a mobility of electrons. It is proportional to the inverse value refined frequency of space-charge waves, defined by [6]

$$\omega_{wsc} = \omega_{pe} - \frac{\omega_{pe}^2}{12 \pi \sigma_d} i \quad (3)$$

where ω_{pe} is frequency of Langmuir oscillations and σ_d is differential conductivity. Between the regions of the electric domain is always set a strong electric field even for small voltages due to the smallness distance between the regions [7]. The origin of the electric domains in the plasma occurs due to the inequality of electron and ion directional drift flows and is accompanied by the generation of transverse electromagnetic waves [6]. The expression for the frequency of these waves can be written as

$$\omega_{shf}^2 = k^2 c^2 + \left(\omega_{pe} - \frac{\omega_{pe}^2}{12 \pi \sigma_d} i \right)^2 \quad (4)$$

The presence of a strong field in the domain and the generation of a transverse electromagnetic wave during its origin allows to accelerate the particles located between the layers of the domain, and particles near it – as a result of their capture by electromagnetic wave. The displacement of a small group of particles at a distance greater than the Debye screening length leads to a significant potential difference between the layers of the electrical domain. Electrons and ions are formed of the neutral atoms due to the high electric field with intensity exceeding the limit value for the appearance of collision ionization.

To comply with a phase-matching condition requires that the particle velocity is close to the phase velocity of electromagnetic waves. Value of voltage on the domain are sufficient to ensure that the electrons have a velocity. The particles accelerated in the area of domain by field receive an additional

acceleration by means of the transverse electromagnetic wave which generated at the separation of charges. The equations of motion in general form for the charged particles in a homogeneous field of a plane electromagnetic wave are [8]

$$\frac{d(m_{\alpha} \vec{u}_{\alpha})}{dt} = q_{\alpha} \left(\vec{E}_z + \frac{\vec{u}_{\alpha} \times \vec{B}_z}{c} \right) \quad (5)$$

in which: α - a sort of particles, q - its charge, \vec{E}_z and \vec{B}_y - components of the electric and magnetic fields ($\vec{E}_z = E_0 e^{i(\alpha x + kz)}$, $\vec{B}_y = B_0 e^{i(\alpha x + ky)}$). The charged particles in the field of electromagnetic wave acquire an energy, that is significantly higher than the value of that corresponds to the applied voltage. The value of acquired energy depends on the values of intensity and induction of fields and time of particles interaction with the wave. Electrical domain performs functions of built-in plasma of microaccelerator of charged particles. Received by the author at the breakdown near surface of a dielectric images of luminosity of plasma are given in Figure 1.




Fig.1. Optical image of luminosity in an initial stage of breakdown -a and the qualitative structure generated under origin of the domain of a transversal electromagnetic waves (show only intensity of electrical field)- b. and luminosity in near-electrode area with secondary V-type formation - c. Designations: A-anode, K-cathode, k-wave vector.

The acceleration of charged particles in plasma on base of electric domain is more efficient than the acceleration on base of laser-plasma interaction. This is due to very low efficiency output of lasers.

References

- [1] A. G. Oreshko, "On domain structures, fields and processes in radiation stars and Universe," in Proc. XXVIII Int. Conf. on Phenomena in Ionized Gases, Prague, 2007, pp. 1888-1891
- [2] J. B. Gunn, "Microwave oscillations of current in III-V semiconductors", Sol. St. Comm, vol. 1, no. 4, pp. 88-91, 1963.
- [3] G. D. Mylnikov, A. P. Napartovich, "The domain instability of a glow discharge", Plasma Physics (Rus. Journal), vol. 1, no. 6, pp. 892-900, 1975.
- [4] A. G. Oreshko, "Double electrical layers of space charge in an cathode plasma," Plasma Physics (Rus. Journal), vol. 17, no. 6, pp. 679-684, 1991.
- [5] M.E.Levinshstein, Yu.K.Pozhela and M.S.Shur *The effect of Gunn* 1975 Moscow: Soviet Radio.
- [6] A. G. Oreshko, "Generation of laboratory ball lightning", Journ. of Physics: Conference series, vol. 44, pp. 127-32, 2006.
- [7] A. G. Oreshko, "The generation of strong fields in plasma," Doklady Physics, vol. 46, pp. 9-11, 2001.
- [8] N.A.Krall, A.W.Trivelpiece *Principle of plasma physics* 1973 NY: McGraw-Hill Book Company.